

isibc/ms/2005/35

May 19th, 2005

<http://www.isibang.ac.in/~statmath/eprints>

On Nyman, Beurling and Baez-Duarte's Hilbert space reformulation of the Riemann Hypothesis

BHASKAR BAGCHI

Indian Statistical Institute, Bangalore Centre
8th Mile Mysore Road-560 059, India

On Nyman, Beurling and Baez-Duarte's Hilbert space reformulation of the Riemann Hypothesis.

Bhaskar Bagchi
Indian Statistical Institute,
Bangalore

Abstract

There has been a surge of interest of late in an old result of Nyman and Beurling giving a Hilbert space formulation of the Riemann Hypothesis. Many authors have contributed to this circle of ideas, culminating in a beautiful refinement due to Baez-Duarte. The purpose of this little survey is to dis-entangle the resulting web of complications, and reveal the essential simplicity of the main results.

Let \mathcal{H} denote the weighted l^2 -space consisting of all sequences $a = \{a_n : n \in \mathbb{N}\}$ of complex numbers such that $\sum_{n=1}^{\infty} \frac{|a_n|^2}{n(n+1)} < \infty$. For any two vectors $a, b \in \mathcal{H}$, their inner product is given by: $\langle a, b \rangle = \sum_{n=1}^{\infty} \frac{a_n \bar{b}_n}{n(n+1)}$. Notice that all bounded sequences of complex numbers are vectors in this Hilbert space. For $l = 1, 2, 3, \dots$ let $\gamma_l \in \mathcal{H}$ be the sequence .

$$\gamma_l = \left\{ \left\{ \frac{n}{l} \right\} : n = 1, 2, 3, \dots \right\}.$$

(Here $\{x\}$, and in what follows, $\{x\}$ is the fractional part of a real number x .) Also, let $\gamma \in \mathcal{H}$ denote the constant sequence

$$\gamma = \{1, 1, 1, \dots\}.$$

Recall that a set A of vectors in a Hilbert space \mathcal{H} is said to be **total** if the set of all finite linear combinations of elements of A is dense in \mathcal{H} , i.e., if no proper closed subspace of the Hilbert space contains the set A . In terms of these few notions and notations, the recent result of Baez-Duarte from [2] can be given the following dramatic formulation.

Theorem 1 *The following statements are equivalent :*

- (i) *The Riemann Hypothesis,*
- (ii) *γ belongs to the closed linear span of $\{\gamma_l : l = 1, 2, 3, \dots\}$, and*
- (iii) *the set $\{\gamma_l : l = 1, 2, 3, \dots\}$ is total in \mathcal{H} .*

We hasten to add that this is not the statement that the reader will see in Baez-Duarte's paper. For one thing, the implications $(ii) \implies (iii)$ and $(iii) \implies (i)$ are not mentioned in this paper : perhaps the author thinks of them as 'well known to experts'. (In such contexts, an expert is usually defined to be a person who has the relevant piece of information.) More over, the main result in [2] is not the

implication (i) \implies (ii) itself, but a ‘unitarily equivalent’ version there-of. More precisely, the result actually proved in [2] is the implication (i) \implies (ii) of Theorem 7 below. In fact, we could not locate in the existing literature the statement (iii) of Theorem 1 (equivalently, of Theorem 7) as a reformulation of the Riemann Hypothesis. This result may be new. It reveals the Riemann Hypothesis as a version of the central theme of Harmonic Analysis : that more or less arbitrary sequences (subject to mild growth restrictions) can be arbitrarily well approximated by superpositions of a class of simple periodic sequences (in this instance, the sequences γ_n).

A second point worth noting is that the particular weight sequence $\{\frac{1}{n(n+1)}\}$ used above is not crucial for the validity of Theorem 1 (though this is the sequence which occurs naturally in its proof). Indeed, any weight sequence $\{w_n : n = 1, 2, 3, \dots\}$ satisfying $\frac{c_1}{n^2} \leq w_n \leq \frac{c_2}{n^2}$ for all n (for constants $0 < c_1 \leq c_2$) would serve equally well. This is because the identity map is an invertible linear operator (hence carrying total sets to total sets) between any two of these weighted l^2 -spaces.

In what follows, we shall adopt the standard practice (in analytic number theory) of denoting a complex variable by $s = \sigma + it$. Thus σ and t are the real and imaginary parts of the complex number s . Recall that **Riemann’s Zeta function** is the analytic function defined on the half-plane $\{\sigma > 1\}$ by the absolutely convergent series $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$. The completed Zeta function ζ^* is defined on this half plane by $\zeta^*(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$, where Γ is Euler’s Gamma function. As Riemann discovered, ζ^* has a meromorphic continuation to the entire complex plane with only two (simple) poles : at $s = 0$ and at $s = 1$. Further, it satisfies the functional equation $\zeta^*(1-s) = \zeta^*(s)$ for all s . Since Γ has poles at the non-positive integers (and nowhere else), it follows that ζ has trivial zeros at the negative even integers. Further, since ζ is real-valued on the real line, its zeros occur in conjugate pairs. This trivial observation, along with the (highly non-trivial) functional equation, shows that the non-trivial zeros of the Zeta function are symmetrically situated about the so-called **critical line** $\{\sigma = \frac{1}{2}\}$. **The Riemann hypothesis** (RH) conjectures that all these non-trivial zeros actually lie on the critical line. In view of the symmetry mentioned above, this amounts to the conjecture that ζ has no zeros on the half-plane

$$\Omega = \{s = \sigma + it : \sigma > \frac{1}{2}, -\infty < t < \infty\}.$$

In other words, the Riemann hypothesis is the statement that $\frac{1}{\zeta}$ is analytic on the half-plane Ω . This is the formulation of RH that we use in this article. Throughout this article, Ω stands for the half-plane $\{\sigma > \frac{1}{2}\}$.

Baez-Duarte’s theorem refines an earlier result of the same type (Theorem 5 below) proved independently by Nyman and Beurling (cf. [6] and [1]). Our intention in this article is to point out that the entire gamut of these results is best seen inside the **Hardy space** $H^2(\Omega)$. Recall that this is the Hilbert space of all analytic functions F on Ω such that

$$\|F\|^2 := \sup_{\sigma > \frac{1}{2}} \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(\sigma + it)|^2 dt < \infty,$$

It is known that any $F \in H^2(\Omega)$ has, almost everywhere, on the critical line, a non-tangential boundary value F^* such that

$$\|F\|^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| F^*\left(\frac{1}{2} + it\right) \right|^2 dt$$

Thus $H^2(\Omega)$ may be identified (via the isometric embedding $F \mapsto F^*$) with a closed subspace of the L^2 -space of the critical line with respect to the Lebesgue measure scaled by the factor $\frac{1}{2\pi}$. (This scaling is to ensure that the Mellin transform F , defined while proving Theorem 2 below, is an isometry.)

For $0 \leq \lambda \leq 1$, let $F_\lambda \in H^2(\Omega)$ be defined by

$$F_\lambda(s) = (\lambda^s - \lambda) \frac{\zeta(s)}{s}, \quad s \in \Omega.$$

Notice that the zero of the first factor at $s = 1$ cancels the pole of the second factor, so that F_λ , thus defined, is analytic on Ω . Also, in view of the well-known elementary estimate (cf. [7])

$$\zeta(s) = O(|s|^{\frac{1}{6}} \log |s|), \quad s \in \overline{\Omega}, \quad s \longrightarrow \infty,$$

the factor $\frac{1}{s}$ ensures that $F_\lambda \in H^2(\Omega)$ for $0 \leq \lambda \leq 1$. (Note that, in order to arrive at this conclusion, any exponent $< \frac{1}{2}$ in the above Zeta estimate would have sufficed. But the exponent $\frac{1}{6}$ happens to be the simplest non-trivial estimate which occurs in the theory of the Riemann Zeta function.) Indeed, under Riemann Hypothesis we have the stronger estimate (Lindelof Hypothesis)

$$\zeta(s) = O(|s|^\epsilon) \text{ as } |s| \longrightarrow \infty, \text{ uniformly for } s \in \overline{\Omega}, \quad (1)$$

for each $\epsilon > 0$. (More precisely, under RH, this estimate holds uniformly on the complement of any given neighbourhood of 1 in $\overline{\Omega}$.)

Finally, for $l = 1, 2, 3, \dots$, let $G_l \in H^2(\Omega)$ be defined by $G_l = F_{\frac{1}{l}}$. Thus,

$$G_l(s) = (l^{-s} - l^{-1}) \frac{\zeta(s)}{s}, \quad s \in \Omega.$$

Also, let $E \in H^2(\Omega)$ be defined by :

$$E(s) = \frac{1}{s}, \quad s \in \Omega.$$

In terms of these notations, the most natural formulation of the Nyman–Beurling–Baez-Duarte theorem is the following :

Theorem 2 *The following statements are equivalent :*

- (i) *The Riemann Hypothesis,*
- (ii) *E belongs to the closed linear span of the set $\{G_l : l = 1, 2, 3, \dots\}$, and*
- (iii) *E belongs to the closed linear span of the set $\{F_\lambda : 0 \leq \lambda \leq 1\}$.*

The plan of the proof is to verify $(i) \implies (ii) \implies (iii) \implies (i)$. As we shall see in a little while, except for the first implication $((i) \implies (ii))$, all these implications are fairly straight forward. In order to prove $(i) \implies (ii)$, we need recall that on the half-plane $\{\sigma > 1\}$, $\frac{1}{\zeta}$ is represented by an absolutely convergent Dirichlet series

$$\sum_{l=1}^{\infty} \mu(l) l^{-s} = \frac{1}{\zeta(s)}. \quad (2)$$

Here $\mu(\cdot)$ is the Mobius function. (To determine its formula, we may formally multiply this Dirichlet series by that of $\zeta(s)$ and equate coefficients to get the recurrence relation $\sum_{l|n} \mu(l) = \delta_{1n}$. Solving this, one can show that $\mu(\cdot)$ takes values in $\{0, +1, -1\}$ and hence the Dirichlet series for $\frac{1}{\zeta}$ is absolutely

convergent on $\{\sigma > 1\}$. Indeed, $\mu(l)$ is $= 0$ if l has a repeated prime factor, is $= +1$ if l has an even number of distinct prime factors, and is $= -1$ if l has an odd number of distinct prime factors. But, for our limited purposes, all this is unnecessary.) What we need is an old theorem of Littlewood (cf. [7]) to the effect that for the validity of the Riemann Hypothesis, it is necessary (and sufficient) that the Dirichlet series displayed above converges uniformly on compact subsets of Ω . Actually, we need the following quantitative version of this theorem of Littlewood.

Lemma 3 *If the Riemann Hypothesis holds then for each $\epsilon > 0$ and each $\delta > 0$, we have $\sum_{l=1}^L \mu(l)l^{-s} = O((|t|+1)^\delta)$ uniformly for $L = 1, 2, 3, \dots$ and uniformly for $s = \sigma + it$ in the half-plane $\{\sigma > \frac{1}{2} + \epsilon\}$. (Thus the implied constant depends only on ϵ and δ .)*

This Lemma may be proved by a minor variation in the original proof of Littlewood's Theorem quoted above. (Note that, with the aid of a little 'normal family' argument, Littlewood's Theorem itself is an easy consequence of this Lemma.) However, for the sake of completeness, we sketch a proof here :

Proof of Lemma 3: We may assume that $s = \sigma + it$ with $\frac{1}{2} + \epsilon < \sigma \leq 1$. (The case $\sigma \geq 1$ is much easier to handle, and we leave out the details.) Fix a positive integer L , and put $x = L + \frac{1}{2}$. Also put $c = 1 - \sigma + \frac{1}{\log x}$. For any large $T > 0$, using residue calculus one can show that for all positive integers n , we have :

$$\frac{1}{2\pi i} \int_{c-iT}^{c+iT} \left(\frac{x}{n}\right)^w \frac{dw}{w} = \begin{cases} 1 + O\left(\frac{(x/n)^c}{T \log(x/n)}\right) & \text{if } n < x, \\ O\left(\frac{(x/n)^c}{T \log(n/x)}\right) & \text{if } n > x. \end{cases}$$

Multiplying this formula by $\mu(n)n^{-s}$ and adding over all positive integers n , we get :

$$\sum_{n=1}^L \mu(n)n^{-s} = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{x^w}{\zeta(s+w)} \frac{dw}{w} + O\left(x^{1-\sigma} \frac{\log(xT)}{T}\right),$$

which is an effective version of Perron's formula. Now, letting $\tilde{c} = \frac{1}{2} + \frac{\delta}{2} - \sigma$, Cauchy's fundamental Theorem yields :

$$\sum_{n=1}^L \mu(n)n^{-s} = \frac{1}{2\pi i} \left(\int_{\tilde{c}-iT}^{\tilde{c}+iT} + \int_{\tilde{c}+iT}^{c+iT} + \int_{c-iT}^{\tilde{c}-iT} \right) \frac{x^w}{\zeta(s+w)} \frac{dw}{w} + \frac{1}{\zeta(s)} + O\left(x^{1-\sigma} \frac{\log(xT)}{T}\right).$$

Now, under RH, we have the wellknown estimate (cf. Theorem 14.2 in [7])

$$\zeta(s)^{-1} = O((|t| + 1)^\epsilon) \tag{3}$$

uniformly for s in the half-plane $\{\sigma \geq \frac{1}{2} + \delta\}$. Therefore the second and third integrals are

$$O\left(x^{1-\sigma} \left(\frac{T^\epsilon + (|t| + 1)^\epsilon}{T}\right)\right),$$

while the first integral is

$$O\left(x^{\frac{1}{2} + \frac{\delta}{2} - \sigma} \log T (T^\epsilon + (|t| + 1)^\epsilon)\right) = O\left(x^{-\delta/2} \log T (T^\epsilon + (|t| + 1)^\epsilon)\right).$$

Combining these estimates and choosing $T = x^B$ where B is a sufficiently small positive constant, we get the required result. ■

Proof of Theorem 2 : (i) \Rightarrow (ii). Assume RH. For positive integers L and any small real number $\epsilon > 0$, let $H_{L,\epsilon} \in H^2(\Omega)$ be defined by

$$H_{L,\epsilon} = \sum_{l=1}^L \frac{\mu(l)}{l^\epsilon} G_l.$$

Thus each $H_{L,\epsilon}$ is in the linear span of $\{G_l : l \geq 1\}$. Note that

$$H_{L,\epsilon}(s) = \frac{\zeta(s)}{s} \left(\sum_{l=1}^L \frac{\mu(l)}{l^{s+\epsilon}} - \sum_{l=1}^L \frac{\mu(l)}{l^{1+\epsilon}} \right), \quad s \in \overline{\Omega}.$$

Therefore, by the Theorem of Littlewood quoted above, for any fixed $\epsilon > 0$,

$$H_{L,\epsilon}(s) \longrightarrow H_\epsilon(s) \quad \text{for } s \text{ in the critical line, as } L \longrightarrow \infty.$$

Here,

$$H_\epsilon(s) := \frac{\zeta(s)}{s} \left(\frac{1}{\zeta(s+\epsilon)} - \frac{1}{\zeta(1+\epsilon)} \right).$$

Also, by the estimates (1), (3) and Lemma 3, $H_{L,\epsilon}$ is bounded by an absolutely square integrable function (viz. a constant times $s^{2\delta-1}$, for any fixed δ in the range $0 < \delta < \frac{1}{4}$). Therefore, by Lebesgue's dominated convergence theorem, we have, for each fixed $\epsilon > 0$,

$$H_{L,\epsilon} \longrightarrow H_\epsilon \quad \text{in the norm of } H^2(\Omega) \text{ as } L \longrightarrow \infty.$$

Since $H_{L,\epsilon}$ is in the linear span of $\{G_l : l = 1, 2, 3, \dots\}$, it follows that, for each $\epsilon > 0$, H_ϵ is in the closed linear span of $\{G_l : l = 1, 2, 3, \dots\}$. Now note that, since ζ has a pole at $s = 1$,

$$H_\epsilon(s) \longrightarrow \frac{1}{s} = E(s) \quad \text{for } s \text{ in the critical line, as } \epsilon \searrow 0.$$

Therefore, in order to show that E is in the closed linear span of $\{G_l : l = 1, 2, 3, \dots\}$ and thus complete this part of the proof, it suffices to show that H_ϵ , $0 < \epsilon < \frac{1}{2}$, are uniformly bounded in modulus on the critical line by an absolutely square integrable function. Then, another application of Lebesgue's dominated convergence would yield

$$H_\epsilon \longrightarrow E \quad \text{in the norm of } H^2(\Omega) \text{ as } \epsilon \searrow 0.$$

Consider the entire function $\xi(s) := s(1-s)\zeta^*(s) = s(1-s)\pi^{-s/2}\Gamma(\frac{s}{2})\zeta(s)$. It has the Hadamard factorisation

$$\xi(s) = \xi(0) \prod_{\rho} \left(1 - \frac{s}{\rho}\right),$$

where the product is over all the non-trivial zeros ρ of the Riemann Zeta function. This product converges provided the zeros ρ and $1-\rho$ are grouped together. In consequence, with a similar bracketing, we have

$$|\xi(s)| = |\xi(0)| \prod_{\rho} \left|1 - \frac{s}{\rho}\right|.$$

Now, under RH, each ρ has real part $= \frac{1}{2}$. Therefore, for s in the closed half-plane $\overline{\Omega}$, we have $|1 - \frac{s}{\rho}| \leq |1 - \frac{s+\epsilon}{\rho}|$. Multiplying this trivial inequality over all ρ , we get

$$|\xi(s)| \leq |\xi(s+\epsilon)|, \quad s \in \overline{\Omega}, \epsilon > 0.$$

(Aside : conversely, the above inequality clearly implies RH. Thus, this simple looking inequality is a reformulation of RH.) In other words, we have, for $s \in \overline{\Omega}$,

$$\left| \frac{\zeta(s)}{\zeta(s+\epsilon)} \right| \leq \pi^{-\epsilon/2} \left| \frac{(s+\epsilon)(1-\epsilon-s)}{s(1-s)} \right| \left| \frac{\Gamma((s+\epsilon)/2)}{\Gamma(s/2)} \right| \leq c \left| \frac{\Gamma((s+\epsilon)/2)}{\Gamma(s/2)} \right|$$

for some absolute constant $c > 0$. But, by Sterling's formula (see [5] for instance), the Gamma ratio on the extreme right is bounded by a constant times $|s|^{\epsilon/2}$, uniformly for $s \in \overline{\Omega}$. Therefore we get

$$\left| \frac{\zeta(s)}{\zeta(s+\epsilon)} \right| \leq c|s|^{\epsilon/2}, \quad s \in \overline{\Omega},$$

for some other absolute constant $c > 0$. In conjunction with the estimate (1), this implies

$$|H_\epsilon(s)| \leq c|s|^{-3/4}, \quad s \in \overline{\Omega},$$

for $0 < \epsilon < \frac{1}{2}$. Since $s \mapsto c|s|^{-3/4}$ is square integrable on the critical line, we are done. This proves the implication (i) \Rightarrow (ii).

Since $\{G_l : l = 1, 2, 3, \dots\} \subseteq \{F_\lambda : 0 \leq \lambda \leq 1\}$, the implication (ii) \Rightarrow (iii) is trivial. To prove (iii) \Rightarrow (i), suppose RH is false. Then there is a Zeta-zero $\rho \in \Omega$. Since $\zeta(\rho) = 0$, it follows that $F_\lambda(\rho) = 0$ for all $\lambda \in (0, 1]$. Thus the set $\{F_\lambda : \lambda \in (0, 1]\}$ (and hence also its closed linear span) is contained in the proper closed subspace $\{F \in H^2(\Omega) : F(\rho) = 0\}$ of $H^2(\Omega)$. (It is a closed subspace since evaluation at any fixed $\rho \in \Omega$ is a continuous linear functional : $H^2(\Omega)$ is a functional Hilbert space.) Since E belongs to the closed linear span of this set, it follows that $0 = E(\rho) = \frac{1}{\rho}$. Hence $0 = 1$: the ultimate contradiction! This proves (iii) \Rightarrow (i). ■

Remark 4 Since $\mu(l) = 0$ unless l is square-free, the functions $H_{L,\epsilon}$ introduced in the course of the above proof are in the linear span of the set $\{G_l : l \text{ square-free}\}$. Thus, the proof actually shows that RH implies (and hence is equivalent to) that E belongs to the closed linear span of the thinner set $\{G_l : l \text{ square-free}\}$ in $H^2(\Omega)$.

Now let $L^2((0, 1])$ be the Hilbert space of complex-valued absolutely square integrable functions (modulo almost everywhere equality) on the interval $(0, 1]$. For $0 \leq \lambda \leq 1$, let $f_\lambda \in L^2((0, 1])$ be defined by

$$f_\lambda(x) = \left\{ \frac{\lambda}{x} \right\} - \lambda \left\{ \frac{1}{x} \right\}, \quad x \in (0, 1].$$

(Recall that $\{\cdot\}$ stands for the fractional part.) Let $\mathbf{1} \in L^2((0, 1])$ denote the constant function $= 1$ on $(0, 1]$. Thus,

$$\mathbf{1}(x) = 1, \quad x \in (0, 1].$$

In terms of these notations, the original theorem of Nyman and Beurling may be stated as :

Theorem 5 *The following statements are equivalent:*

- (i) *The Riemann Hypothesis,*
- (ii) *$\mathbf{1}$ is in the closed linear span in $L^2((0, 1])$ of the set $\{f_\lambda : 0 \leq \lambda \leq 1\}$,*
- (iii) *the set $\{f_\lambda : 0 \leq \lambda \leq 1\}$ is total in $L^2((0, 1])$.*

Proof : One defines the *Fourier-Mellin transform* $F : L^2((0, 1]) \longrightarrow H^2(\Omega)$ by :

$$F(f)(s) = \int_0^{\infty} x^{s-1} f(x) dx, \quad s \in \Omega, \quad f \in L^2((0, 1]). \quad (4)$$

It is wellknown that F , thus defined, is an isometry. For completeness, we sketch a proof. Since $s \longmapsto (x \longmapsto x^{s-1})$ is an $L^2((0, 1])$ -valued analytic function on Ω , it follows that $F(f)$ is analytic on Ω for each $f \in L^2((0, 1])$. For $\lambda \in [0, 1]$, let $\Psi_\lambda \in L^2((0, 1])$ denote the indicator function of the interval $(0, \lambda)$. Using the well-known identity

$$\frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{e^{iux}}{1+x^2} dx = e^{-|u|}, \quad u \in \mathbb{R},$$

one sees that $\|F(\Psi_\lambda)\|^2 = \|\Psi_\lambda\|^2 < \infty$ – hence $F(\Psi_\lambda) \in H^2(\Omega)$ – and, more generally, $\|F(\Psi_\lambda) - F(\Psi_\mu)\|^2 = \|\Psi_\lambda - \Psi_\mu\|^2$ for $\lambda, \mu \in [0, 1]$. Since $\{\Psi_\lambda : \lambda \in [0, 1]\}$ is a total subset of $L^2((0, 1])$, this implies that F maps $L^2((0, 1])$ isometrically into $H^2(\Omega)$.

We begin with a computation of the Melin transform of f_λ . Claim :

$$F(f_\lambda) = -F_\lambda, \quad 0 \leq \lambda \leq 1. \quad (5)$$

To verify this claim, begin with $s = \sigma + it$, $\sigma > 1$. Then, $\int_0^1 \left\{\frac{\lambda}{x}\right\} x^{s-1} dx = \lambda \int_0^1 x^{s-2} dx - \int_0^1 \left[\frac{\lambda}{x}\right] x^{s-1} dx = \frac{\lambda}{s-1} - \int_0^1 \left[\frac{\lambda}{x}\right] x^{s-1} dx$. But,

$$\begin{aligned} \int_0^1 \left[\frac{\lambda}{x}\right] x^{s-1} dx &= \sum_{n=1}^{\infty} n \int_{\lambda/(n+1)}^{\lambda/n} x^{s-1} dx \\ &= \frac{\lambda^s}{s} \sum_{n=1}^{\infty} n(n^{-s} - (n+1)^{-s}). \end{aligned}$$

Now, the partial sum $\sum_{n=1}^N n(n^{-s} - (n+1)^{-s})$ telescopes to $-N(N+1)^{-s} + \sum_{n=1}^N n^{-s}$. Since $\sigma > 1$, letting $N \longrightarrow \infty$, we get $\sum_{n=1}^{\infty} n(n^{-s} - (n+1)^{-s}) = \zeta(s)$. Thus,

$$\int_0^1 \left\{\frac{\lambda}{x}\right\} x^{s-1} dx = \frac{\lambda}{s-1} - \lambda^s \frac{\zeta(s)}{s}.$$

In particular, taking $\lambda = 1$ here, one gets

$$\int_0^1 \left\{\frac{1}{x}\right\} x^{s-1} dx = \frac{1}{s-1} - \frac{\zeta(s)}{s}.$$

Multiplying the second equation by λ and subtracting the result from the first, we arrive at

$$\int_0^1 f_\lambda(x) x^{s-1} dx = -(\lambda^s - \lambda) \frac{\zeta(s)}{s} = -F_\lambda(s)$$

for s in the half-plane $\{\sigma > 1\}$. Since both sides of this equation are analytic in the bigger half-plane Ω , this equation continues to hold for $s \in \Omega$. This proves the Claim (??).

(i) \implies (ii). Assume RH. Then, by Theorem 2, $E = F(\mathbf{1})$ belongs to the closed linear span of $\{F_\lambda = -F(f_\lambda) : 0 \leq \lambda \leq 1\}$. Since F is an isometry, this shows that $\mathbf{1}$ belongs to the closed linear span of the set $\{f_\lambda : 0 \leq \lambda \leq 1\}$. Thus (i) \implies (ii).

(ii) \implies (iii). Let $\mathbf{1}$ be in the closed linear span in $L^2((0, 1])$ of $\{f_\lambda : 0 \leq \lambda \leq 1\}$. Applying F , it follows that E is in the closed linear span (say \mathcal{N}) of $\{F_\lambda : 0 \leq \lambda \leq 1\}$. For $\mu \in (0, 1]$, let $\Theta_\mu \in H^\infty(\Omega)$ (the Banach algebra of bounded analytic functions on Ω) be defined by

$$\Theta_\mu(s) = \mu^{s-\frac{1}{2}}, \quad s \in \Omega.$$

We have $|\Theta_\mu(s)| = 1$ for s in the critical line. That is, Θ_μ is an inner function. In consequence, the linear operators $M_\mu : H^2(\Omega) \longrightarrow H^2(\Omega)$ defined by

$$M_\mu(F) = \Theta_\mu F \quad (\text{point-wise product}), \quad F \in H^2(\Omega),$$

are isometries. (Since $\Theta_\lambda \Theta_\mu = \Theta_{\lambda\mu}$, it follows that $M_\lambda M_\mu = M_{\lambda\mu}$ for $\lambda, \mu \in (0, 1]$. Thus $\{M_\mu : \mu \in (0, 1]\}$ is a semi-group of isometries on $H^2(\Omega)$ modelled after the multiplicative semi-group $(0, 1]$.) Trivially, for $0 \leq \lambda \leq 1$ and $0 < \mu \leq 1$, we have:

$$M_\mu(F_\lambda) = \Theta_\mu F_\lambda = \mu^{-1/2}(F_{\lambda\mu} - \lambda F_\mu).$$

This shows that the closed subspace \mathcal{N} spanned by the F_λ 's is invariant under the semi-group $\{M_\mu : \mu \in (0, 1]\}$:

$$M_\mu(\mathcal{N}) \subseteq \mathcal{N}, \quad \mu \in (0, 1].$$

Since $E \in \mathcal{N}$, it follows that $M_\mu(E) \in \mathcal{N}$ for $\mu \in (0, 1]$. But we have the trivial computation

$$F(\Psi_\lambda) = \lambda^{1/2} M_\lambda(E), \quad 0 < \lambda \leq 1.$$

Thus, $\{F(\Psi_\lambda) : 0 \leq \lambda \leq 1\}$ is contained in the closed linear span \mathcal{N} of $\{F(f_\lambda) : 0 \leq \lambda \leq 1\}$. Since F is an isometry, it follows that $\{\Psi_\lambda : 0 \leq \lambda \leq 1\}$ is contained in the closed linear span in $L^2((0, 1])$ of the set $\{f_\lambda : 0 \leq \lambda \leq 1\}$. Since the first set is clearly total in $L^2((0, 1])$, it follows that so is the second. Thus (ii) \implies (iii).

(iii) \implies (i). Clearly (iii) implies that the closed linear span of $\{f_\lambda : 0 \leq \lambda \leq 1\}$ contains $\mathbf{1}$ and hence, applying F , the closed linear span of $\{F_\lambda : 0 \leq \lambda \leq 1\}$ contains E . Therefore, by Theorem 2, Riemann Hypothesis follows. Thus (iii) \implies (i). ■

Remark 6 *It is instructive to compare the proof of Theorem 5 with Beurling's original proof as given in [4]. Our proof makes it clear that the heart of the matter is very simple : Riemann Hypothesis amounts to the existence of approximate inverses to the Zeta function in a suitable function space (viz. the weighted Hardy space of analytic functions on Ω with the weight function $|E(s)|^2$). The simplification in its proof is achieved by Baez-Duarte's perfectly natural and yet vastly illuminating observation that, under RH, these approximate inverses are provided by the partial sums of the Dirichlet series for $\frac{1}{\zeta}$. In contrast, Beurling's original proof is a clever and ill-motivated application of Phragmen-Lindelof type arguments. (We have not seen Nyman's original proof.) To be fair, we should however point out that such arguments are now hidden under the carpet : they occur in the proofs (not presented here) of the conditional estimates (3) and (1).*

Let \mathcal{M} be the closed subspace of $L^2((0, 1])$ consisting of the functions which are almost everywhere constant on each of the sub-intervals $(\frac{1}{n+1}, \frac{1}{n}]$, $n = 1, 2, 3, \dots$. Since each element of \mathcal{M} is almost everywhere equal to a unique function which is everywhere constant on these sub-intervals, we may (and do) think of \mathcal{M} as the space of all such (genuine) piece-wise constant functions. As a closed subspace of a Hilbert space, \mathcal{M} is a Hilbert space in its own right.

For $l = 1, 2, 3, \dots$, let $g_l \in L^2((0, 1])$ be defined by

$$g_l(x) = \left\{ \frac{1}{lx} \right\} - \frac{1}{l} \left\{ \frac{1}{x} \right\}, \quad x \in (0, 1].$$

Thus, $g_l = f_{1/l}$, $l = 1, 2, 3, \dots$

Notice that we have $g_l(x) = \frac{1}{l} \left[\frac{1}{x} \right] - \left[\frac{1}{lx} \right]$. Also, for $x \in (\frac{1}{n+1}, \frac{1}{n}]$, $n = 1, 2, 3, \dots$, $\frac{1}{lx} \in [\frac{n}{l}, \frac{n+1}{l})$, and no integer can be in the interior of the latter interval, so that $\left[\frac{1}{lx} \right] = \left[\frac{n}{l} \right]$; also, $\left[\frac{1}{x} \right] = n$ for $x \in (\frac{1}{n+1}, \frac{1}{n}]$. Thus we get:

$$g_l(x) = g_l\left(\frac{1}{n}\right) = \left\{ \frac{n}{l} \right\}, \quad x \in \left(\frac{1}{n+1}, \frac{1}{n} \right]. \quad (6)$$

In consequence,

$$g_l \in \mathcal{M}, \quad l = 1, 2, 3, \dots$$

The refinement of Baez-Duarte of the Beurling-Nyman theorem may now be stated as follows. (However, as already stated, the implication (i) \implies (ii) of this theorem is its only part which explicitly occurs in [2].)

Theorem 7 *The following are equivalent :*

- (i) *The Riemann Hypothesis,*
- (ii) *$\mathbf{1}$ belongs to the closed linear span of $\{g_l : l = 1, 2, 3, \dots\}$, and*
- (iii) *$\{g_l : l = 1, 2, 3, \dots\}$ is a total set in \mathcal{M} .*

Proof : Putting $\lambda = \frac{1}{l}$ in the Formula (??), we get :

$$F(g_l) = -G_l, \quad l = 1, 2, 3, \dots$$

Since, under RH, $E = F(\mathbf{1})$ is in the closed linear span of $\{G_l = -F(g_l) : l = 1, 2, 3, \dots\}$ and F is an isometry, it follows that $\mathbf{1}$ is in the closed linear span of $\{g_l : l = 1, 2, 3, \dots\}$. Thus (i) \implies (ii).

Now, for positive integers m , define the linear operators $T_m : \mathcal{M} \longrightarrow \mathcal{M}$ by :

$$(T_m f)(x) = \begin{cases} m^{1/2} f(mx) & \text{if } x \in (0, \frac{1}{m}], \\ 0 & \text{if } x \in (\frac{1}{m}, 1]. \end{cases}$$

Clearly each T_m is an isometry. (We have $T_m T_n = T_{mn}$ – thus $\{T_m : m = 1, 2, 3, \dots\}$ is a semigroup of isometries modelled after the multiplicative semi-group of positive integers.) Also, it is easy to see that

$$T_m(g_l) = m^{1/2} (g_{lm} - \frac{g_m}{l})$$

for any two positive integers l, m . Thus the closed linear span \mathcal{K} of the vectors g_l , $l = 1, 2, 3, \dots$ is invariant under this semi-group. Further, letting $\Phi_n \in \mathcal{M}$ denote the indicator function of the interval $(0, \frac{1}{n}]$, one has :

$$T_m(\Phi_n) = m^{1/2} \Phi_{mn}.$$

Thus, if \mathcal{K} contains $\mathbf{1} = \Phi_1$ then it contains Φ_n for all n . Since $\{\Phi_n : n = 1, 2, 3, \dots\}$ is clearly a total subset of \mathcal{M} , it then follows that $\mathcal{K} = \mathcal{M}$, so that $\{g_l : l = 1, 2, 3, \dots\}$ is a total subset of \mathcal{M} . Thus (ii) \implies (iii).

Lastly, if $\{g_l : l = 1, 2, 3, \dots\}$ is a total subset of \mathcal{M} then, in particular its closed linear span contains $\mathbf{1}$, and hence the closed linear span of $\{G_l = -F(g_l)\}$ contains $E = F(\mathbf{1})$, so that RH follows by Theorem 2. Thus (iii) \implies (i). ■

Proof of Theorem 1: Let $U : \mathcal{M} \longrightarrow \mathcal{H}$ be the unitary defined by

$$U(f) = \left\{ f\left(\frac{1}{n}\right) : n = 1, 2, 3, \dots \right\}, \quad f \in \mathcal{M}.$$

Since $U(\mathbf{1}) = \gamma$ and (in view of the Formula (??)) $U(g_l) = \gamma_l$, this Theorem is a straightforward reformulation of Theorem 7. ■

Remark 8 *In view of Remark 4, Riemann hypothesis actually implies (and hence is equivalent to) the statement that γ belongs to the closed linear span in \mathcal{H} of the much thinner set $\{\gamma_l : l \text{ square-free}\}$.*

So where does the undoubtedly elegant reformulation of RH in Theorem 1 leave us? One possible approach is as follows. For positive integers L , let $D(L)$ denote the distance of the vector $\gamma \in \mathcal{H}$ from the $(L - 1)$ -dimensional subspace of \mathcal{H} spanned by $\gamma_1, \gamma_2, \dots, \gamma_L$. In view of Theorem 1, RH is equivalent to the statement $D(L) \longrightarrow 0$ as $L \longrightarrow \infty$. So one might try to estimate $D(L)$. Indeed, as a discrete analogue of a conjecture of Baez-Duarte et. al. in [3], one might expect that $D^2(L)$ is asymptotically equal to $\frac{A}{\log L}$ for $A = 2 + C - \log(4\pi)$, where C is Euler's constant. (But, of course, this is far stronger than RH itself.) A standard formula gives $D^2(L)$ as a ratio of two Gram determinants, i.e., determinants with the inner products $\langle \gamma_l, \gamma_m \rangle$ as entries. It is easy to write down these inner products as finite sums involving the logarithmic derivative of the Gamma function. But such formulae are hardly suitable for calculation/estimation of determinants. In any case, it will be a sad day for Mathematics when (and if) the Riemann Hypothesis is proved by a brute-force calculation! Surely a dramatically new and deep idea is called for. But then, as a wise man once said, it is fool-hardy to predict – specially the future!

References

- [1] A. Beurling, A closure problem related to the Riemann Zeta function, Proc. Nat. Acad. Sci. 41, 1955, 312-314
- [2] L Baez-Duarte, A strengthening of the Nyman–Beurling criterion for the Riemann Hypothesis, Atti Acad. Naz. Lincei 14, 2003, 5–11
- [3] L. Baez-Duarte, M. Balazard, B. Landreau and E. Saias, Notes sur la fonction ζ de Riemann 3, Advances in Math. 149, 2000, 130-144.
- [4] W.F. Donoghue, jr., Distributions and Fourier Transforms, Academic Press, 1969.
- [5] S. Lang, Complex Analysis, Springer Verlag, 1992
- [6] B. Nyman, On some groups and semi-groups of translations, Ph.D. Thesis, Uppsala, 1950.
- [7] E.C. Titchmarsh, The theory of the Riemann Zeta function, Oxford Univ. Press, 1951.